
BOUNDS FOR p -ADIC EXPONENTIAL SUMS AND LOG-CANONICAL THRESHOLDS

by

Raf Cluckers & Willem Veys

Abstract. — We propose a conjecture for exponential sums which generalizes both a conjecture by Igusa and a local variant by Denef and Sperber, in particular dropping the homogeneity condition on the polynomial f , and with new uniform properties. The exponential sums have summation set consisting of integers modulo p^m lying p -adically close to y , and the proposed bounds are uniform in p , y , and m . We give some evidence for the conjecture, by showing uniform bounds in p , y and small values for m . Our method consists in bounding the log-canonical threshold in relation to the bounds predicted by the conjecture, and reducing to finite field exponential sums about which we use results by Katz.

1. Introduction and main results

Let us fix a nonconstant polynomial F in n variables over \mathbb{Z} . We consider, for any integer $m > 1$ and any prime number p , the global exponential sum

$$S(F, p, m) := p^{-mn} \cdot \sum_{x \in (\mathbb{Z}/p^m\mathbb{Z})^n} \exp(2\pi i \frac{F(x)}{p^m}),$$

2000 Mathematics Subject Classification. — Primary 11L07; Secondary 11L05.

Key words and phrases. — p -adic exponential sums, Denef-Sperber conjecture, Igusa's conjecture on exponential sums, log-canonical threshold, motivic oscillation index.

and, for any $y \in \mathbb{Z}^n$, its local version

$$S_y(F, p, m) := p^{-mn} \cdot \sum_{x \in y + (p\mathbb{Z}/p^m\mathbb{Z})^n} \exp(2\pi i \frac{F(x)}{p^m}),$$

where

$$y + (p\mathbb{Z}/p^m\mathbb{Z})^n = \{x \in (\mathbb{Z}/p^m\mathbb{Z})^n \mid x_i \equiv y_i \pmod{p} \text{ for each } i\}.$$

Based on somewhat less general conjectures of Igusa [11] and of Denef and Sperber [8], we conjecture bounds for the complex modulus of the above sums, which are uniform in p, m , and y . We express these bounds in terms of the log-canonical thresholds, but a stronger formulation in terms of e.g. the motivic oscillation index of [4] would also make sense. For any field k of characteristic zero, a polynomial $f \in k[x] = k[x_1, \dots, x_n]$ and a point $y \in k^n$ satisfying $f(y) = 0$, we write $c_y(f)$ to denote the log-canonical threshold of f at y (see Definition 2.1 below), and $c(f)$ for the log canonical threshold of f , being the minimum of the $c_y(f)$ when y runs over all points in \bar{k}^n satisfying $f(y) = 0$, where \bar{k} is an algebraic closure of k .

Let us first fix some more notation.

Definition 1.1. — Let $a(F)$ be the minimum, over all $b \in \mathbb{C}$, of the log-canonical thresholds of the polynomials $F(x) - b$. Further, for $y \in \mathbb{Z}^n$, let $a_{y,p}(F)$ be the minimum of the log-canonical thresholds at y' of the polynomials $F(x) - F(y')$, where the minimum is taken over all $y' \in y + (p\mathbb{Z}_p)^n$. Note that $a(F) \leq a_{y,p}(F)$ for each p and y .

Conjecture 1.2. — *There exists a function $L_F : \mathbb{N} \rightarrow \mathbb{N}$ with $L_F(m) \ll m^{n-1}$ such that for all primes p , all $m \geq 2$, and all $y \in \mathbb{Z}^n$, one has*

$$(1.2.1) \quad |S(F, p, m)|_{\mathbb{C}} \leq L_F(m) p^{-ma(F)}$$

and

$$(1.2.2) \quad |S_y(F, p, m)|_{\mathbb{C}} \leq L_F(m) p^{-ma_{y,p}(F)},$$

where $|\cdot|_{\mathbb{C}}$ is the complex norm.

Under some extra conditions that were introduced by Igusa for reasons of adelic integrability, but that we believe irrelevant for bounding the above sums, he conjectured in the introduction of [11] that (1.2.1) holds for all homogeneous F and all $m \geq 1$. We believe that focusing on m at least 2 allows one to remove the homogeneity condition, and we give evidence below. The bounds (1.2.1) (with the log-canonical threshold,

resp. the variant with the motivic oscillation index in the exponent) imply Igusa's original conjecture (with the log-canonical threshold, resp. any of his proposed candidate oscillation indices in the exponent), including the case $m = 1$, by [4]. Indeed, the case $m = 1$ of Igusa's conjecture is known by [4] for any of these exponents. The estimates (1.2.1) of the conjecture yield a criterion to show adelic L^q -integrability for an adelic function related to $S(F, p, m)$, with a simple lower bound on q based on the exponent $a(F)$, as noted by Igusa in [11]. Denef and Sperber [8] conjectured the local variant (1.2.2) for $y = 0$, thus without uniformity in y . Both inequalities, namely the global (1.2.1) and the local but uniform (1.2.2), seem closely related.

We prove Conjecture 1.2 for m up to the value 4, and, more generally, for m up to some value related to orders of vanishing, defined as follows.

Definition 1.3. — Let r be the minimum of the order of vanishing of the functions $x \mapsto F(x) - b$ at the singular points in \mathbb{C}^n of $F = b$, i.e., the minimum of the multiplicities of the singular points of the hypersurfaces $F = b$, where b runs over \mathbb{C} . Here we consider the minimum over the empty set to be $+\infty$. Further, for $y \in \mathbb{Z}^n$, let $r_{y,p}$ be the minimum of the order of vanishing of the functions $x \mapsto F(x) - F(y')$ at y' , where y' runs only over singular points in the p -adic neighbourhood $y + (p\mathbb{Z}_p)^n$ for which moreover $c_{y'}(F - F(y')) = a_{y,p}(F)$.

Note that by definition $r \geq 2$ and $r_{y,p} \geq 2$. With notation as introduced above and with $+\infty + a = +\infty$ for any real a , we can now state our main result as evidence for Conjecture 1.2.

Theorem 1.4. — *There exists a constant L_F such that, for all prime numbers p , all $y \in \mathbb{Z}^n$, and all m with $2 \leq m \leq r + 2$, resp. with $2 \leq m \leq r_{y,p} + 2$, one has*

$$(1.4.1) \quad |S(F, p, m)|_{\mathbb{C}} \leq L_F p^{-ma(F)},$$

resp.

$$(1.4.2) \quad |S_y(F, p, m)|_{\mathbb{C}} \leq L_F p^{-ma_{y,p}(F)}.$$

In Section 2.6, we explain analogues over finite field extensions of \mathbb{Q}_p and over $\mathbb{F}_q((t))$, for large primes p and q any power of p .

Theorem 1.4 is proved using new inequalities for log-canonical thresholds and by reducing to finite field exponential sums for which bounds by Katz can be used, see Lemma 2.3.

Let us now explain the bounds on log-canonical thresholds related to the conjecture. Let f be a nonconstant polynomial over \mathbb{C} in the variables $x = (x_1, \dots, x_n)$, and write

$$(1.4.3) \quad f = \sum_{i \geq r} f_i,$$

with f_i either identically zero or homogeneous and of degree i , and where f_r is nonzero for some $r \geq 2$. As before, write $c_0(f)$ for the log-canonical threshold of f at zero. If f is non-reduced at zero (that is, g^2 divides f for some polynomial g which vanishes at 0), then one knows that

$$(1.4.4) \quad c_0(f) \leq \frac{1}{2}.$$

In any case one has (see Section 8 of [13])

$$(1.4.5) \quad c_0(f) \leq \frac{n}{r}.$$

The following inequalities can be considered as a certain combination of the above two (quite obvious) inequalities, but with the non-reducedness assumption on f_r , instead of on f .

Lemma 1.5. — *Suppose that g^2 divides f_r for some nonconstant polynomial g . Then one has the inequality*

$$(1.5.1) \quad (r+1)c_0(f) \leq n + \frac{1}{2}.$$

If moreover g divides f_{r+1} (this includes the case f_{r+1} identically zero), then

$$(1.5.2) \quad (r+2)c_0(f) \leq n + 1.$$

Lemma 1.5 will be obtained as a corollary of the following sharper and unconditional bounds, which we think are of independent interest.

Proposition 1.6. — *With notation from (1.4.3), one has*

$$(1.6.1) \quad (r+1)c_0(f) \leq n + c(f_r).$$

One should compare (1.6.1) with the bound $|c_0(f) - c(f_r)| \leq n/(r+1)$ from Proposition 8.19 of [13]. A generalization of Proposition 1.6, with a bound for $(e+1)c_0(f)$ for arbitrary $e > 0$, is given at the end of the paper, see Theorem 2.10. By combining Lemma 1.5 with results from [10], we obtain global variants.

Proposition 1.7. — *Let $r > 1$ be an integer and let f be a polynomial in n variables and with complex coefficients. Suppose that, for y running over an irreducible d -dimensional variety $Y \subset \mathbb{C}^n$, one has that f vanishes with order at least r at y . For $y \in Y$, let us write $f_y(x)$ for the polynomial $f(x+y)$ in the variables x , and $f_y = \sum_{i \geq r} f_{y,i}$ with $f_{y,i}$ either identically zero or homogeneous and of degree i . Then one has*

$$(1.7.1) \quad rc_y(f) \leq n - d$$

and, for a generic $y \in Y$,

$$(1.7.2) \quad (r+1)c_y(f) \leq n - d + c(f_{y,r}).$$

In particular, for a generic $y \in Y$, if $f_{y,r}$ is non-reduced, then

$$(1.7.3) \quad (r+1)c_y(f) \leq n - d + \frac{1}{2}.$$

If, for a generic $y \in Y$, there is a non-constant polynomial g_y which divides $f_{y,r+1}$ and such that g_y^2 divides $f_{y,r}$, then one further has

$$(1.7.4) \quad (r+2)c_y(f) \leq n - d + 1.$$

The proofs of Theorem 1.4, Proposition 1.6, Lemma 1.5 and the global variants are given in Section 2.

1.8. Some context and notation. — Conjecture 1.2 is known when the implied constant is allowed to depend on the prime number p , see [11] and [9]. Namely, for each prime p there exists a function $L_{F,p} : \mathbb{N} \rightarrow \mathbb{N}$ with $L_{F,p}(m) \ll m^{n-1}$, such that for all $m \geq 2$ and all $y \in \mathbb{Z}_p^n$, both estimates

$$(1.8.1) \quad |S(F, p, m)|_{\mathbb{C}} \leq L_{F,p}(m)p^{-ma(F)}$$

and

$$(1.8.2) \quad |S_y(F, p, m)|_{\mathbb{C}} \leq L_{F,p}(m)p^{-ma_{y,p}(F)}$$

hold. In the case that F is non-degenerate with respect to (the compact faces of) the Newton polyhedron at zero of F , then the bounds (1.2.2) with $y = 0$ hold, see [8] and [5]. If F is non-degenerate and quasi-homogeneous, then also the bounds from (1.2.1) hold, by [8] and [5]. For other work on Igusa's original conjecture, we refer to [3], [4], [14], [16]. Lemma 5.4 of [2] gives other evidence for Conjecture 1.2, under some specific geometric conditions. Related exponential sums in few variables (namely with small n) have been studied in [14], [16] and in [6], [7].

Below we will write $|\cdot|$ instead of $|\cdot|_{\mathbb{C}}$ for the complex norm. For complex valued functions H and G on a set Z , the notation $H \ll G$ means that there exists a constant $c > 0$ such that $|H(z)| \leq c|G(z)|$ for all z in Z .

All integrals over K^n , for any non-archimedean local field K with valuation ring \mathcal{O}_K , will be against the Haar measure $|dx|$ on K^n , normalized so that \mathcal{O}_K^n has measure 1.

We write $\mathbb{F}_p^{\text{alg}}$ for an algebraic closure of \mathbb{F}_p .

2. Proofs of the main results

We first recall two descriptions of the log-canonical threshold.

Definition 2.1. — For a non-constant polynomial f in n variables over an algebraically closed field K of characteristic zero, and $y \in K^n$ satisfying $f(y) = 0$, the log-canonical threshold of f at y is denoted by $c_y(f)$ and defined as follows. For any proper birational morphism $\pi : Y \rightarrow K^n$ from a smooth variety Y , and for any prime divisor E on Y , we denote by N and $\nu - 1$ the multiplicities along E of the divisors of π^*f and $\pi^*(dx_1 \wedge \cdots \wedge dx_n)$, respectively. Then

$$c_y(f) = \inf_{\pi, E} \left\{ \frac{\nu}{N} \right\},$$

where π runs over all π as above and E over all prime divisors on Y such that $y \in \pi(E)$. For a polynomial f over a non-algebraically closed field k of characteristic zero and $y \in k^n$ satisfying $f(y) = 0$, one defines $c_y(f)$ as above with K any algebraic closure of k . Finally, when f is the zero polynomial, one defines $c(f)$ as 0.

In fact $c_y(f) = \min_E \left\{ \frac{\nu}{N} \right\}$, where π is any fixed embedded resolution of the germ of $f = 0$ at y (and $y \in \pi(E)$). Note that always $c_y(f) \leq 1$, a property not shared by the motivic oscillation index of f .

By Mustașă's Corollaries 0.2 and 3.6 of [15], we can describe the log-canonical threshold by taking certain dimensions, as follows.

Let p be an integer and h a nonconstant polynomial over \mathbb{C} in n variables. Write $\text{Cont}^{\geq p}(h)$ for the subset of $\mathbb{C}[[t]]^n$ given by

$$\{x \in \mathbb{C}[[t]]^n \mid h(x) \equiv 0 \pmod{(t^p)}\}$$

and $\text{Cont}_0^{\geq p}(h)$ for

$$\{x \in \mathbb{C}[[t]]^n \mid \text{ord}_t h(x) \equiv 0 \pmod{(t^p)}, x \in (t\mathbb{C}[[t]])^n\}.$$

Let us further write

$$\text{codim Cont}^{\geq p}(h)$$

for the codimension of $\rho_m(\text{Cont}^{\geq p}(h))$ in $\rho_m(\mathbb{C}[[t]]^n)$ for any $m \geq p$, where $\rho_m : \mathbb{C}[[t]]^n \rightarrow (\mathbb{C}[t]/(t^{m+1}))^n$ is the projection modulo t^{m+1} in each coordinate. Here, $\rho_m(\text{Cont}^{\geq p}(h))$ is seen as a Zariski closed subset of $\mathbb{C}^{n(m+1)} \cong \rho_m(\mathbb{C}[[t]]^n)$. The definition is independent of the choice of m . We write similarly $\text{codim Cont}_0^{\geq p}(h)$ for the codimension of $\rho_m(\text{Cont}_0^{\geq p}(h))$ in $\rho_m(\mathbb{C}[[t]]^n)$ for any $m \geq p$.

By Corollary 0.2 of [15], for all integers $k > 0$, we have

$$(2.1.1) \quad c(h) \leq \frac{\text{codim Cont}^{\geq k}(h)}{k}$$

and there exist infinitely many $k > 0$ for which equality holds. Also, if h vanishes at 0, one has by Corollary 3.6 of [15] that

$$(2.1.2) \quad c_0(h) = \inf_{k>0} \frac{\text{codim Cont}_0^{\geq k}(h)}{k}.$$

Based on these relations, we can now prove Proposition 1.6.

Proof of Proposition 1.6. — By the equality statement concerning (2.1.1) for f_r , there exists $k > 0$ such that

$$(2.1.3) \quad c(f_r) = \frac{\text{codim Cont}^{\geq k}(f_r)}{k}.$$

Let ℓ be $kr + k$. Now define the cylinder $B \subset \mathbb{C}[[t]]^n$ as

$$B := \{x \in \mathbb{C}[[t]]^n \mid \rho_{k-1}(x) = \{0\}, \text{ord}_t f_r(x) \geq \ell\}.$$

By the homogeneity of f_r , the cylinder B can be considered (under corresponding identifications), as

$$\rho_{k-1}(B) \times t^k \text{Cont}^{\geq k}(f_r) = \{0\} \times t^k \text{Cont}^{\geq k}(f_r) \subset \mathbb{C}[[t]]^n.$$

Again by the homogeneity of f_r and the fact that $f - f_r$ has multiplicity at least $r + 1$, one has

$$B \subset \text{Cont}_0^{\geq \ell}(f).$$

Hence, by (2.1.2), one finds

$$(2.1.4) \quad c_0(f) \leq \frac{\text{codim } B}{\ell},$$

where $\text{codim } B$ is defined as the codimension of $\rho_m(B)$ in $\rho_m(\mathbb{C}[[t]]^n)$ for large enough m . On the other hand, one finds from (2.1.1) that

$$\text{codim } B = kn + \text{codim}(\text{Cont}^{\geq k}(f_r)) = kn + kc(f_r).$$

Using this together with (2.1.4) and dividing by k , one finds (1.6.1). \square

It is also possible to give a proof for Proposition 1.6 based on embedded resolution of singularities, without using Mustaă's formulas.

Alternative proof of Proposition 1.6. — Let $\pi_0 : Y_0 \rightarrow \mathbb{C}^n$ be the blowing-up at the origin; its exceptional divisor E_0 is projective $(n-1)$ -space. We consider for example the chart on Y_0 where E_0 is given by $x_1 = 0$ and π_0^*f by

$$x_1^r \left(f_r(1, x_2, \dots, x_n) + x_1 \sum_{i \geq r+1} x_1^{i-r-1} f_i(1, x_2, \dots, x_n) \right).$$

Along E_0 the multiplicity of the pullback of $dx = dx_1 \wedge \dots \wedge dx_n$ is n and the multiplicities of both π_0^*f and $\pi_0^*f_r$ are r .

We now perform a composition of blowing-ups $Y \rightarrow Y_0$, leading to an embedded resolution $\pi : Y \rightarrow \mathbb{C}^n$ of $f_r = 0$. More precisely, for example on the chart above, we only use centres ‘not involving x_1 ’; hence they all have positive dimension and are transversal to E_0 . Say $c(f_r) = \frac{\nu}{N}$, where E is an exceptional component of π such that along E the multiplicities of the pullback of dx and f_r are ν and N , respectively. We may assume that $E \neq E_0$; otherwise $c(f_r) = \frac{n}{r}$ and the statement becomes trivial.

Consider analytic or étale coordinates x_1, y_2, \dots, y_n in a generic point of $E \cap E_0 \subset Y$ such that E is given by $y_2 = 0$. In that point π^*f is of the form

$$x_1^r \left(y_2^N u(y_2, \dots, y_n) + x_1(\dots) \right),$$

where $u(y_2, \dots, y_n)$ is a unit. Next, we blow up Y at the codimension two centre $Z_1 = E \cap E_0$ given (locally) by $x_1 = y_2 = 0$. Along the new exceptional divisor E_1 the multiplicities of the pullback of dx and f are $n + \nu$ and $r + \mu_1$, respectively, where $\mu_1 \geq 1$ is the order of vanishing of $y_2^N u(y_2, \dots, y_n) + x_1(\dots)$, the strict transform of f , along Z_1 . In fact, in the relevant chart the pullback of f is now of the form

$$x_1^r y_2^{r+\mu_1} \left(y_2^{N-\mu_1} u(y_2, \dots, y_n) + x_1(\dots) \right).$$

As long as E_0 intersects the strict transform of $f = 0$, we continue to blow up with centre this intersection, in the relevant chart always given

by $x_1 = y_2 = 0$. Let E_k be the last exceptional component created this way. Then along E_k the multiplicities of the pullback of dx and f are $kn + \nu$ and $kr + \sum_{i=1}^k \mu_i$, respectively, where the μ_i are the orders of vanishing of the strict transform of f along the centres of blow-up. Note that $\sum_{i=1}^k \mu_i = N$. We just showed that

$$(2.1.5) \quad c_0(f) \leq \frac{kn + \nu}{kr + N}.$$

An elementary computation, using that $\frac{\nu}{N} \leq \frac{n}{r}$ and $k \leq N$, shows that

$$(2.1.6) \quad \frac{kn + \nu}{kr + N} \leq \frac{n + \frac{\nu}{N}}{r + 1} = \frac{n + c(f_r)}{r + 1}.$$

Then combining (2.1.5) and (2.1.6) finishes the proof. \square

Remark 2.2. — (1) The proof above can be shortened by using a weighted blow-up instead of the last k blow-ups.

(2) M. Mustașă informed us of yet another proof of Proposition 1.6, using multiplier ideals.

Proof of Lemma 1.5. — The inequality (1.5.1) follows from (1.6.1) and (1.4.4) for f_r . For inequality (1.5.2) and with g as in the lemma, consider the cylinder C given by

$$\{x \in \mathbb{C}[[t]]^n \mid \rho_0(x) = 0, \text{ord}_t g(\frac{x_1}{t}, \dots, \frac{x_n}{t}) \geq 1\}.$$

Then one easily verifies that

$$C \subset \text{Cont}_0^{\geq r+2}(f)$$

and $\text{codim } C = n + 1$. The result now follows from Mustașă's bound as in (2.1.2) for f and $k = r + 2$. \square

Proof of Proposition 1.7. — By Theorem 1.2 of [10], one has for generic y in Y and a generic vector subspace H of \mathbb{C}^n of dimension $n - d$ that

$$c_0(f_{y|H}) = c_y(f),$$

where $f_{y|H}$ is the restriction of the polynomial map f_y to H . The proposition now follows from the genericity of y and H , by (1.4.5) and by Proposition 1.6 and Lemma 1.5 applied to $f_{y|H}$. \square

In the proof of our main theorems we will use the following lemmas. The first one follows almost directly from work by Katz in [12] and Noether normalization.

Lemma 2.3. — *Let n, k, N be nonnegative integers. Then there exist constants D and E such that the following hold for all prime numbers p with $p > E$, all positive powers q of p , and all nontrivial additive characters ψ_q on \mathbb{F}_q . Let g_1, \dots, g_k and h be (nonconstant) homogeneous polynomials in $x = (x_1, \dots, x_n)$ with coefficients in \mathbb{Z} and of degree at most N . Let X be the reduced subscheme of $\mathbb{A}_{\mathbb{Z}}^n$ associated to the ideal (g_1, \dots, g_k) .*

If h (modulo p) does not vanish on any irreducible component of $X_p := X \otimes \mathbb{F}_p^{\text{alg}}$ of dimension equal to $\dim X_p$, then

$$(2.3.1) \quad \left| \sum_{y \in X(\mathbb{F}_q)} \psi_q(h(y)) \right| \leq D \cdot q^{\dim X_p - 1/2}.$$

If the image of h in $\mathbb{F}_p^{\text{alg}}[x]$ under $\mathbb{Z}[x] \rightarrow \mathbb{F}_p^{\text{alg}}[x]$ is reduced, then

$$(2.3.2) \quad \left| \sum_{y \in \mathbb{F}_q^n} \psi_q(h(y)) \right| \leq D \cdot q^{n-1}.$$

Proof. — The bounds in (2.3.2) follow immediately from Katz [12], Theorem 4. In the case that X_p is irreducible, the bounds in (2.3.1) follow from Theorem 5 of [12]. The remaining case that X_p is reducible follows from the irreducible case and Noether normalization. \square

From now on, let F and r be as in the introduction. We will use some instances of the Ax-Kochen principle of [1], like the following lemma.

Lemma 2.4. — *For large enough p , any $v \in \mathbb{F}_p^n$, and any $y \in \mathbb{Z}_p^n$ lying above v , the following holds. If the reduction of F modulo p vanishes with order r at v , then*

$$\text{ord}(F(y)) \geq r,$$

where ord is the p -adic order $\mathbb{Q}_p \rightarrow \mathbb{Z} \cup \{+\infty\}$.

Proof. — The statement is easily reduced to a simple statement over a discrete valuation ring of equicharacteristic zero. One finishes by a standard ultraproduct argument (namely by the Ax-Kochen principle). \square

Lemma 2.5. — *Let V be the subscheme of $\mathbb{A}_{\mathbb{Z}}^n$ given by the equations $\text{grad } F = 0$. If p is large enough, then one has for any $m > 1$ that*

$$S(F, p, m) = \sum_{v \in V(\mathbb{F}_p)} \int_{u \in \mathbb{Z}_p^n, u \equiv v \pmod{p}} \exp(2\pi i \frac{F(u)}{p^m}) |du|$$

and that $S_y(F, p, m) = 0$ whenever the reduction of y modulo p does not lie in $V(\mathbb{F}_p)$.

Proof. — This follows by Hensel's Lemma and by the basic relation

$$\sum_{t \in \mathbb{F}_p} \psi_p(t) = 0$$

for any nontrivial additive character ψ_p on \mathbb{F}_p . \square

We begin with the proof of the almost trivial part of Theorem 1.4.

Proof of Theorem 1.4 for $m \leq r$, resp. $m \leq r_{y,p}$. — Note that for small p , there is nothing to prove by (1.8.1), resp. (1.8.2). If $r = +\infty$, the theorem follows easily. We may thus suppose that $r < +\infty$ and that p is large. Let V be the subscheme of $\mathbb{A}_{\mathbb{Z}}^n$ given by the equations $\text{grad } F = 0$, and write d for the dimension of $V \otimes \mathbb{C}$. Fix $m > 1$ with $m \leq r$, resp. $m \leq r_{y,p}$. For all $y \in \mathbb{Z}^n$ one has

$$ma(F) \leq ra(F) \leq n - d,$$

by (1.7.1), resp.

$$ma_{y,p}(F) \leq r_{y,p}a_{y,p}(F).$$

Also, when p is large enough, one has

$$S(F, p, m) = p^{-n} \#V(\mathbb{F}_p),$$

resp.,

$$(2.5.1) \quad S_y(F, p, m) = p^{-n} \text{ and } r_{y,p}a_{y,p}(F) \leq n$$

for $y \bmod p$ in $V(\mathbb{F}_p)$, and

$$S_y(F, p, m) = 0$$

for $y \bmod p$ outside $V(\mathbb{F}_p)$. Indeed, this follows by Lemmas 2.4 and 2.5. By Noether normalization there exists D such that

$$\#V(\mathbb{F}_p) \leq Dp^d,$$

uniformly in p . One readily finds

$$|S(F, p, m)| \leq Dp^{-ma(F)},$$

resp.

$$|S_y(F, p, m)| \leq p^{-ma_{y,p}(F)},$$

for all large p and all $y \in \mathbb{Z}^n$, which finishes the proof. \square

Proof of Theorem 1.4 for $m = r + 1$, resp. $m = r_{y,p} + 1$

Note that for small p , there is nothing to prove by (1.8.1), resp. (1.8.2). We may thus again suppose that p is large and that $r < +\infty$. Fix $y \in \mathbb{Z}^n$. By Lemma 2.5 we may suppose that there exists a critical point $y' \in y + \mathbb{Z}_p^n$ of F , such that $F - F(y')$ vanishes with order $r_{y,p}$ at y' and $c_{y'}(F - F(y')) = a_{y,p}(F)$. Write $f_y(x)$ for $F(x + y') - F(y')$ and $f_y = \sum_{i \geq r_{y,p}} f_{y,i}$ with $f_{y,i}$ either identically zero or homogeneous and of degree i and with $f_{y,r_{y,p}}$ nonzero for a choice of such y' . We first prove (1.4.2) by the following calculation, where ψ is the additive character on \mathbb{Q}_p sending x to $\exp(2\pi i x')$ for any rational number x' which lies in $\mathbb{Z}[1/p]$ and satisfying $x - x' \in \mathbb{Z}_p$, and with Haar measure normalized as in section 1.8:

$$\begin{aligned}
S_y(F, p, r_{y,p} + 1) &= \int_{x \in y + (p\mathbb{Z}_p)^n} \psi\left(\frac{F(x)}{p^{r_{y,p}+1}}\right) |dx| \\
&= \int_{x \in (p\mathbb{Z}_p)^n} \psi\left(\frac{f_y(x) + F(y')}{p^{r_{y,p}+1}}\right) |dx| \\
&= \frac{b_y}{p^n} \int_{u \in \mathbb{Z}_p^n} \psi\left(\frac{p^{r_{y,p}} f_{y,r_{y,p}}(u) + p^{r_{y,p}+1} f_{y,r_{y,p}+1}(u) + \dots}{p^{r_{y,p}+1}}\right) |du| \\
&= \frac{b_y}{p^n} \int_{u \in \mathbb{Z}_p^n} \psi\left(\frac{p^{r_{y,p}} f_{y,r_{y,p}}(u)}{p^{r_{y,p}+1}}\right) |du| \\
&= \frac{b_y}{p^n} \int_{u \in \mathbb{Z}_p^n} \psi\left(\frac{f_{y,r_{y,p}}(u)}{p}\right) |du| \\
&= \frac{b_y}{p^n} \sum_{v \in \mathbb{F}_p^n} \int_{u \in \mathbb{Z}_p^n, \bar{u}=v} \psi\left(\frac{f_{y,r_{y,p}}(u)}{p}\right) |du| \\
&= \frac{b_y}{p^{2n}} \sum_{v \in \mathbb{F}_p^n} \psi_p(\overline{f_{y,r_{y,p}}}(v)).
\end{aligned}$$

Here we denote by \bar{u} the tuple in \mathbb{F}_p^n obtained by reduction mod p of the components $u_i \in \mathbb{Z}_p$ of u , by ψ_p the nontrivial additive character on \mathbb{F}_p sending w to $\psi(w'/p)$ for any $w' \in \mathbb{Z}_p$ which projects to w , by $\overline{f_{y,r_{y,p}}}$ the reduction modulo p of $f_{y,r_{y,p}}$, and we put

$$b_y := \psi\left(\frac{F(y')}{p^{r_{y,p}+1}}\right).$$

Now by Lemma 2.3, applied to $h = f_{y,r_{y,p}}$ and with $k = 0$, there exists a constant $D > 0$ such that

$$\left| \sum_{v \in \mathbb{F}_p^n} \psi_p(\overline{f_{y,r_{y,p}}}(v)) \right| \leq D \cdot p^{n-\delta_{y,p}}$$

for each large p and uniformly in y for $\delta_{y,p}$ so that $\delta_{y,p} = 1/2$ in the case that $\overline{f_{y,r_{y,p}}}$ is non-reduced, and $\delta_{y,p} = 1$ in the case that $\overline{f_{y,r_{y,p}}}$ is reduced.

We claim, for large p and for all $y \in \mathbb{Z}^n$, that

$$(2.5.2) \quad (r_{y,p} + 1)c_0(f_y) \leq n + \delta_{y,p}.$$

If y' is a non-isolated critical point of F (in the set of critical points of F with coordinates in an algebraic closure of \mathbb{Q}_p), then $r_{y,p}c_0(f_y) \leq n - 1$ by (1.7.1) and the claim follows from $c_0(f_y) \leq 1$. Also, if $\delta_{y,p} = 1$, then the claim follows from (1.4.5) and $c_0(f_y) \leq 1$. In the case that y' is an isolated critical point (in the set of critical points of F with coordinates in an algebraic closure of \mathbb{Q}_p) and $\delta_{y,p} = 1/2$ simultaneously, it follows from our assumption that p is large that $\overline{f_{y,r_{y,p}}}$ is non-reduced and thus (2.5.2) follows from Lemma 1.5. This assumption of p being large is uniform in y since there are only finitely many isolated critical points of F . Hence, we find for all large p and all y that

$$(2.5.3) \quad |S_y(F, p, r_{y,p} + 1)| = \frac{1}{p^{2n}} \left| \sum_{v \in \mathbb{F}_p^n} \psi_p(\overline{f_{y,r_{y,p}}}(v)) \right|$$

$$\leq D \cdot p^{-n-\delta_{y,p}}$$

$$(2.5.4) \quad \leq D \cdot p^{-(r_{y,p}+1)c_0(f_y)} \leq D \cdot p^{-(r_{y,p}+1)a_{y,p}(F)}.$$

This completes the proof of (1.4.2) for all y and $m = r_{y,p} + 1$.

To show (1.4.1), let V be the subscheme of $\mathbb{A}_{\mathbb{Z}}^n$ given by the equations $\text{grad } F = 0$, and let d be the dimension of $V \otimes \mathbb{C}$. For each $v \in V(\mathbb{F}_p)$, fix a point $y(v)$ in \mathbb{Z}^n lying above v , and a critical point $y'(v)$ of F lying above v such that $F - F(y'(v))$ vanishes with order $r_{y(v),p}$ and $c_{y'}(F - F(y')) = a_{y,p}(F)$ (such y' exists since p is assumed large). Now (1.4.1) for $m = r + 1$ follows by estimating, for large primes p ,

$$(2.5.5) \quad |S(F, p, r + 1)| = \left| \sum_{v \in V(\mathbb{F}_p)} S_{y(v)}(F, p, r + 1) \right|$$

$$(2.5.6) \quad \leq \sum_{v \in V(\mathbb{F}_p)} D \cdot p^{-n-\varepsilon_v},$$

for some $D > 0$, and where ε_v equals $\delta_{y(v),p}(y'(v))$ whenever $r = r_{y,p}$ and where $\varepsilon_v = 0$ when $r < r_{y,p}$. Here the equality (2.5.5) follows from Lemma 2.4, and the inequality (2.5.6) comes from (2.5.3) when $r = r_{y,p}$ and from (2.5.1) when $r < r_{y,p}$. By quantifier elimination for the language of rings with coefficients in \mathbb{Z} , there exist V_0 , $V_{1/2}$, and V_1 , such that V_i is a finite disjoint union of subschemes of V (it is constructible and defined over \mathbb{Z}) with $\cup_i V_i(\mathbb{C}) = V(\mathbb{C})$ and such that the following hold, for $i = 0, \frac{1}{2}$, and 1. The polynomial $F - F(b)$ vanishes with order $> r$ at b for $b \in V_0(\mathbb{C})$, $F - F(b)$ vanishes with order r at b for $b \in V_{1/2}(\mathbb{C})$ and also for $b \in V_1(\mathbb{C})$, and $(F(x+b) - F(b))_r$ is reduced for $b \in V_1(\mathbb{C})$, and non-reduced for $b \in V_{1/2}(\mathbb{C})$. Let d_i be the dimension of $V_i \otimes \mathbb{C}$. Note that for large p , one has $\varepsilon_v = i$ for $v \in V_i(\mathbb{F}_p)$. Now we bound as follows:

$$(2.5.7) \quad |S(F, p, r+1)| \leq \sum_i \#V_i(\mathbb{F}_p) D \cdot p^{-n-i}$$

$$(2.5.8) \quad \leq \sum_i \#V_i(\mathbb{F}_p) \cdot D \cdot p^{-(r+1)a(F)-d_i}$$

$$(2.5.9) \quad \leq D' p^{-ma(F)},$$

for some D' . The inequality (2.5.7) follows from (2.5.6), (2.5.8) follows from Proposition 1.7 and the definition of $a(F)$ as a minimum, and (2.5.9) from Noether normalization. \square

Proof of Theorem 1.4 for $m = r + 2$, resp. $m = r_{y,p} + 2$.

For the same reasons as in the previous proofs we may concentrate on large primes p and suppose $r < +\infty$. Fix $y \in \mathbb{Z}^n$. By Lemma 2.5 we may suppose that there exists a critical point $y' \in y + \mathbb{Z}_p^n$ of F , such that $F - F(y')$ vanishes with order $r_{y,p}$ at y' and $c_{y'}(F - F(y')) = a_{y,p}(F)$. Write $f_y(x)$ for $F(x + y') - F(y')$ and $f_y = \sum_{i \geq r_{y,p}} f_{y,i}$ with $f_{y,i}$ either identically zero or homogeneous and of degree i , and where $f_{y,r_{y,p}}$ is nonzero. We first prove (1.4.2). Let X be the subscheme of $\mathbb{A}_{\mathbb{Z}_p}^n$ associated to the equations $\text{grad } f_{y,r_{y,p}} = 0$. Let A_p be the subset of \mathbb{Z}_p^n of those points whose projection mod p lies in $X(\mathbb{F}_p)$. Also, let C_p be the

complement of A_p in \mathbb{Z}_p^n . We calculate as follows:

$$\begin{aligned}
S_y(F, p, r_{y,p} + 2) &= \int_{x \in y + (p\mathbb{Z}_p)^n} \psi\left(\frac{F(x)}{p^{r_{y,p}+2}}\right) |dx| \\
&= \frac{b_y}{p^n} \int_{u \in \mathbb{Z}_p^n} \psi\left(\frac{p^{r_{y,p}} f_{y,r_{y,p}}(u) + p^{r_{y,p}+1} f_{y,r_{y,p}+1}(u)}{p^{r_{y,p}+2}}\right) |du| \\
&= \frac{b_y}{p^n} \int_{u \in \mathbb{Z}_p^n} \psi\left(\frac{f_{y,r_{y,p}}(u) + p f_{y,r_{y,p}+1}(u)}{p^2}\right) |du| \\
&= \frac{b_y}{p^n} (I_1 + I_2),
\end{aligned}$$

where $b_y = \psi\left(\frac{F(y')}{p^{r_{y,p}+2}}\right)$,

$$I_1 = I_1(y) = \int_{u \in A_p} \psi\left(\frac{f_{y,r_{y,p}}(u) + p f_{y,r_{y,p}+1}(u)}{p^2}\right) |du|,$$

and

$$I_2 = I_2(y) = \int_{u \in C_p} \psi\left(\frac{f_{y,r_{y,p}}(u) + p f_{y,r_{y,p}+1}(u)}{p^2}\right) |du|.$$

One has $I_2 = 0$ by Hensel's Lemma and by the basic relation

$$\sum_{t \in \mathbb{F}_p} \psi_p(t) = 0$$

for the nontrivial additive character ψ_p on \mathbb{F}_p .

To estimate $|I_1|$, we first assume the condition on y and y' that $f_{y,r_{y,p}+1}$ vanishes on at least one absolutely irreducible component of X of maximal dimension. We will show that this condition on y and y' implies

$$(2.5.10) \quad (r_{y,p} + 2)c_0(f_y) \leq 2n - \dim(X \otimes \mathbb{Q}_p).$$

If $\dim(X \otimes \mathbb{Q}_p) \leq n - 2$, then (2.5.10) follows from $(r_{y,p} + 2)c_0(f_y) \leq n + 2$, which in turn follows from $c_0(f_y) \leq 1$ and (1.4.5). If $\dim X \otimes \mathbb{Q}_p = n - 1$ one has that $(r_{y,p} + 2)c_0(f_y) \leq n + 1$ by Lemma 1.5, and (2.5.10) follows also in this case and thus in general. By Noether normalization, there exists $E > 0$ independent of y such that

$$\#X(\mathbb{F}_p) \leq Ep^{\dim(X \otimes \mathbb{Q}_p)}$$

for all large p . Since

$$|I_1| \leq \frac{\#X(\mathbb{F}_p)}{p^n},$$

we find from the above discussion that, for all y satisfying the above condition,

$$\frac{1}{p^n} |I_1| \leq Ep^{\dim(X \otimes \mathbb{Q}_p) - 2n} \leq Ep^{-(r_{y,p}+2)c_0(f_y)} \leq Ep^{-(r_{y,p}+2)a_{y,p}(F)}$$

for all large p .

Finally assume the condition on y and y' that $f_{y,r_{y,p}+1}$ does not vanish on any absolutely irreducible component of X of maximal dimension. By Lemma 2.4, one can rewrite I_1 for large p as

$$I_1 = \int_{u \in A_p} \psi\left(\frac{f_{y,r_{y,p}+1}(u)}{p}\right) |du|.$$

Using this expression we compute

$$\begin{aligned} \frac{1}{p^n} I_1 &= \frac{1}{p^n} \sum_{v \in X(\mathbb{F}_p)} \int_{\bar{u}=v, u \in \mathbb{Z}_p^n} \psi\left(\frac{f_{y,r_{y,p}+1}(u)}{p}\right) |du| \\ &= \frac{1}{p^{2n}} \sum_{v \in X(\mathbb{F}_p)} \psi_p(\overline{f_{y,r_{y,p}+1}}(v)), \end{aligned}$$

where the notations \bar{u} , ψ_p , and $\overline{f_{y,r_{y,p}+1}}$ are as in the proof of the case $m = r_{y,p} + 1$, namely reductions modulo p . By Lemma 2.3, there exists $N > 0$ such that, for all y satisfying the above condition, and for all large p ,

$$\left| \sum_{y \in X(\mathbb{F}_p)} \psi_p(\overline{f_{y,r_{y,p}+1}}(y)) \right| \leq Np^{\dim(X \otimes \mathbb{Q}_p) - 1/2}.$$

Hence,

$$\left| \frac{1}{p^n} I_1 \right| \leq Np^{-2n + \dim(X \otimes \mathbb{Q}_p) - 1/2}$$

for large p . If $f_{y,r_{y,p}}$ is non-reduced, then $\dim X \otimes \mathbb{Q}_p = n - 1$. If $f_{y,r_{y,p}}$ is reduced, then $\dim(X \otimes \mathbb{Q}_p) \leq n - 2$. By (1.5.1) of Lemma 1.5, $c_0(f_y) \leq 1$ and (1.4.5), one finds in any case that

$$(r_{y,p} + 2)c_0(f_y) \leq 2n - \dim(X \otimes \mathbb{Q}_p) + 1/2.$$

Hence,

$$\frac{1}{p^n} |I_1| \leq Np^{-(r_{y,p}+2)c_0(f_y)} \leq Np^{-(r_{y,p}+2)a_{y,p}(F)} = Np^{-ma_{y,p}(F)}$$

for each large p , which finishes the proof of (1.4.2) for $m = r_{y,p} + 2$. One derives (1.4.1) for $m = r + 2$ by adapting the argument showing (1.4.2) as in the proof for $m = r + 1$. \square

2.6. Finite field extensions. — As usual it is possible to prove analogous uniform bounds for all finite field extensions of \mathbb{Q}_p and all fields $\mathbb{F}_q((t))$, when one restricts to large residue field characteristics. We just give the definitions and formulate the analogue of Conjecture 1.2 and the analogue of Theorem 1.4.

Let \mathcal{O} be a ring of integers of a number field, and let $N > 0$ be an integer. Let F be a polynomial with coefficients in $\mathcal{O}[1/N]$ in the variables $x = (x_1, \dots, x_n)$. Let $\mathcal{C}_{\mathcal{O}[1/N]}$ be the collection of all non-archimedean local fields K (of any characteristic) with a ring homomorphism $\mathcal{O}[1/N] \rightarrow K$ (where local means locally compact). For K in $\mathcal{C}_{\mathcal{O}[1/N]}$, write \mathcal{O}_K for its valuation ring with maximal ideal \mathcal{M}_K and residue field k_K with q_K elements. Further write $\psi_K : K \rightarrow \mathbb{C}^\times$ for an additive character which is trivial on the valuation ring \mathcal{O}_K and nontrivial on $\pi_K^{-1}\mathcal{O}_K$ where π_K is a uniformizer of \mathcal{O}_K . The analogue of the above integrals $S(F, p, m)$ and $S_y(F, p, m)$ for K in $\mathcal{C}_{\mathcal{O}[1/N]}$ are the following integrals for λ in K^\times ,

$$S(F, K, \lambda) := \int_{x \in \mathcal{O}_K^n} \psi_K\left(\frac{F(x)}{\lambda}\right) |dx|$$

and, for $y \in \mathcal{O}_K^n$,

$$S_y(F, K, \lambda) := \int_{x \in y + (\mathcal{M}_K)^n} \psi_K\left(\frac{F(x)}{\lambda}\right) |dx|,$$

where $|dx|$ is the Haar measure on K^n , normalized such that \mathcal{O}_K^n has measure one, and where $y + (\mathcal{M}_K)^n = \prod_{i=1}^n (y_i + \mathcal{M}_K)$.

The following naturally generalizes Conjecture 1.2, again formulated with the log-canonical threshold in the exponent, where other exponents, like the motivic oscillation index, that can be larger than 1, also would make sense.

Conjecture 2.7. — *There exist $M > 0$ and a function $L_F : \mathbb{N} \rightarrow \mathbb{N}$ with $L_F(m) \ll m^{n-1}$ such that for all $K \in \mathcal{C}_{\mathcal{O}[1/N]}$ whose residue field has characteristic at least M , all $y \in \mathcal{O}_K^n$, and all $\lambda \in K^\times$ with $\text{ord}(\lambda) \geq 2$, if one writes $m = \text{ord}(\lambda)$, one has*

$$|S(F, K, \lambda)|_{\mathbb{C}} \leq L_F(m) q_K^{-ma(F)},$$

and

$$|S_y(F, K, \lambda)|_{\mathbb{C}} \leq L_F(m) q_K^{-ma_{y,K}(F)}.$$

Here ord denotes the valuation on K^\times with $\text{ord}(\pi_K) = 1$, and $a_{y,K}(F)$ equals the minimum of the log-canonical thresholds of $F(x) - F(y')$ at y' , where the minimum is taken over all $y' \in y + (\mathcal{M}_K)^n$.

With the same proof as for Theorem 1.4, we find the following.

Theorem 2.8. — *Let F be a polynomial over $\mathcal{O}[1/N]$. There exist $M > 0$ and a constant L_F such that for all $K \in \mathcal{C}_{\mathcal{O}[1/N]}$ whose residue field has characteristic at least M and for all $\lambda \in K^\times$, if one writes $m = \text{ord}(\lambda)$ and if $2 \leq m \leq r + 2$, resp. $2 \leq m \leq r_{y,K} + 2$, then one has*

$$|S(F, K, \lambda)|_{\mathbb{C}} \leq L_F q_K^{-ma(F)},$$

resp.

$$|S_y(F, K, \lambda)|_{\mathbb{C}} \leq L_F q_K^{-ma_{y,K}(F)}.$$

Here, $r_{y,K}(F)$ is the minimum of the order of vanishing of $x \mapsto F(x) - F(y')$ where x runs over those singular points of the polynomial mapping $x \mapsto F(x) - F(y') : y + \prod_{i=1}^n (y_i + \mathcal{M}_K) \rightarrow K$ with $c_{y'}(F - F(y')) = a_{y,p}(F)$.

2.9. A recursive bound for $c_0(f)$. — We conclude the paper with a generalization of the bound of Proposition 1.6, which also sharpens (1.5.2). Let f be a non-constant polynomial over \mathbb{C} in the variables $x = (x_1, \dots, x_n)$ with $f(0) = 0$, and write

$$(2.9.1) \quad f = \sum_{i \geq 1} f_i,$$

with f_i either identically zero or homogeneous of degree i .

For e a positive integer, let d_e be the least common multiple of the integers $1, 2, \dots, e$, and let $I_e(f)$ be the ideal generated by the polynomials

$$f_i^{d_e/(e-i+1)}$$

for i with $1 \leq i \leq e$. Write $c(I_e(f))$ for the log-canonical threshold of the ideal $I_e(f)$. (The log canonical threshold $c(I)$ of a non-zero ideal I in n variables over \mathbb{C} can be defined analogously as in Definition 2.1, for instance as $\min_E \{\frac{\nu}{N}\}$, where π is now any fixed log-principalization of I and N is now the multiplicity along E of the divisor of $I\mathcal{O}_Y$. See e.g. [15] for more details.) We put $c(I) = 0$ when I is the zero ideal.

Theorem 2.10. — *One has for any $e > 0$ that*

$$(2.10.1) \quad (e + 1)c_0(f) \leq n + d_e \cdot c(I_e(f)).$$

Before proving Theorem 2.10, we state an equivalent formulation and give some illustrative examples of (2.10.1).

Write as usual $f = \sum_{i \geq r} f_i$, where f_r is nonzero. For k a positive integer, let $J_k(f)$ be the ideal generated by the polynomials

$$f_{r+i}^{d_k/(k-i)}$$

for i with $0 \leq i \leq k-1$. Then

$$(2.10.2) \quad (r+k)c_0(f) \leq n + d_k \cdot c(J_k(f)).$$

This reformulation (2.10.2) follows directly from (2.10.1), using the multiplicativity of the log-canonical threshold, namely, that $a \cdot c(I^a) = c(I)$ for any integer $a > 0$ and any ideal I . Its advantage is that the involved numbers are smaller.

For $k = 1$, we obtain

$$(r+1)c_0(f) \leq n + c(f_r),$$

which is Proposition 1.6. The case $k = 2$ sharpens and generalizes (1.5.2):

$$(r+2)c_0(f) \leq n + 2c(f_r, f_{r+1}^2).$$

As a third example, for $k = 3$, we have

$$(r+3)c_0(f) \leq n + 6c(f_r^2, f_{r+1}^3, f_{r+2}^6).$$

The proof of Theorem 2.10 is similar to the first one of Proposition 1.6.

Proof of Theorem 2.10. — For any ideal I of $\mathbb{C}[x]$ and any integer $p > 0$, we will write $\text{Cont}^{\geq p}(I)$ for

$$\{x \in \mathbb{C}[[t]]^n \mid \text{ord}_t h(x) \equiv 0 \pmod{t^p}, \text{ for all } h \in I\}.$$

By Corollary 3.4 of [15], there exists $k > 0$ such that

$$(2.10.3) \quad d_e k c(I_e(f)) = \text{codim } \text{Cont}^{\geq d_e k}(I_e(f)),$$

where the codimension is taken as before (namely after projecting by ρ_m for high enough m). Now define the cylinder $B \subset \mathbb{C}[[t]]^n$ with $\rho_{k-1}(B) = \rho_{k-1}(\{0\}) = \{0\}$ and, (under corresponding identifications)

$$B := \rho_{k-1}(\{0\}) \times t^k \text{Cont}^{\geq d_e k}(I_e(f)) = \{0\} \times t^k \text{Cont}^{\geq d_e k}(I_e(f)) \subset \mathbb{C}[[t]]^n.$$

By the homogeneity of the f_i , one checks for each i that

$$B \subset \text{Cont}_0^{\geq k(e+1)}(f_i),$$

and we have thus that

$$B \subset \mathrm{Cont}_0^{\geq k(e+1)}(f).$$

Hence, by Corollary 3.6 of [15], one finds

$$(2.10.4) \quad k(e+1)c_0(f) \leq \mathrm{codim} B.$$

On the other hand, one finds by (2.10.3) and the definition of B that

$$\mathrm{codim} B = kn + \mathrm{codim}(\mathrm{Cont}_0^{\geq d_e k}(I_e(f))) = kn + d_e k c(I_e(f)).$$

Using this together with (2.10.4) and dividing by k , one finds (2.10.1). \square

Remark 2.11. — Also for Theorem 2.10, we could give another proof along the lines of the alternative proof of Proposition 1.6. More precisely, one blows up the origin, constructs a log-principalization of the ideal $I_e(f)$, and performs an adequate weighted blow-up in order to obtain an exceptional component with the desired numerical invariants.

Acknowledgments. — The authors would like to thank J. Denef and S. Sperber for their generous sharing of insights. We would like to express special thanks towards M. Mustařă and J. Nicaise for discussions around Proposition 1.6 and Theorem 2.10.

The authors were supported in part by the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) with ERC Grant Agreements nrs. 246903 NMNAG and 615722 MOTMEL-SUM, by the Labex CEMPI (ANR-11-LABX-0007-01), and by the Research Fund KU Leuven (grant OT11/069).

References

- [1] J. Ax and S. Kochen, *Diophantine problems over local fields. I*, Amer. J. Math. **87** (1965), 605–630.
- [2] B. J. Birch, *Forms in many variables*, Proc. Roy. Soc. Ser. A **265** (1961/1962), 245–263. MR 0150129 (27 #132)
- [3] R. Cluckers, *Igusa and Denef-Sperber conjectures on nondegenerate p -adic exponential sums*, Duke Math. J. **141** (2008), no. 1, 205–216.
- [4] ———, *Igusa's conjecture on exponential sums modulo p and p^2 and the motivic oscillation index*, Int. Math. Res. Not. IMRN **2008** (2008), no. 4, article ID rnm118, 20 pages.
- [5] ———, *Exponential sums: questions by Denef, Sperber, and Igusa*, Trans. Amer. Math. Soc. **362** (2010), no. 7, 3745–3756.

- [6] Todd Cochrane and Zhiyong Zheng, *Pure and mixed exponential sums*, Acta Arith. **91** (1999), no. 3, 249–278. MR 1735676 (2000k:11093)
- [7] ———, *Exponential sums with rational function entries*, Acta Arith. **95** (2000), no. 1, 67–95. MR 1787206 (2001g:11130)
- [8] J. Denef and S. Sperber, *Exponential sums mod p^n and Newton polyhedra*, Bull. Belg. Math. Soc. Simon Stevin **suppl.** (2001), 55–63.
- [9] J. Denef and W. Veys, *On the holomorphy conjecture for Igusa’s local zeta function*, Proc. Amer. Math. Soc. **123** (1995), no. 10, 2981–2988.
- [10] A. Dimca, Ph. Maisonobe, M. Saito, and T. Torrelli, *Multiplier ideals, V -filtrations and transversal sections*, Math. Ann. **336** (2006), no. 4, 901–924. MR 2255178 (2007h:32042)
- [11] J. Igusa, *Lectures on forms of higher degree (notes by S. Raghavan)*, Lectures on mathematics and physics, Tata institute of fundamental research, vol. 59, Springer-Verlag, 1978.
- [12] N. Katz, *Estimates for "singular" exponential sums*, Int. Math. Res. Not. IMRN (1999), no. 16, 875–899.
- [13] János Kollár, *Singularities of pairs*, Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 221–287. MR 1492525 (99m:14033)
- [14] B. Lichtin, *On a conjecture of Igusa*, Mathematika **59** (2013), no. 2, 399–425.
- [15] Mircea Mustață, *Singularities of pairs via jet schemes*, J. Amer. Math. Soc. **15** (2002), no. 3, 599–615 (electronic). MR 1896234 (2003b:14005)
- [16] J. Wright, *Exponential sums and polynomial congruences in two variables: the quasi-homogeneous case*, arXiv:1202.2686.

RAF CLUCKERS, Université Lille 1, Laboratoire Painlevé, CNRS - UMR 8524, Cité Scientifique, 59655 Villeneuve d’Ascq Cedex, France, and, Katholieke Universiteit Leuven, Department of Mathematics, Celestijnenlaan 200B, B-3001 Leuven, Belgium, • *E-mail* : Raf.Cluckers@math.univ-lille1.fr
Url : <http://math.univ-lille1.fr/~cluckers>

WILLEM VEYS, KU Leuven, Department of Mathematics, Celestijnenlaan 200B, B-3001 Leuven, Belgium • *E-mail* : wim.veys@wis.kuleuven.be
Url : <http://wis.kuleuven.be/algebra/veys.htm>